



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2005

Adamantanethione

Heimgartner, Heinz ; Mlostoń, Grzegorz ; Romanski, Jaroslaw

DOI: <https://doi.org/10.1002/047084289X.rn00504>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-63513>

Book Section

Accepted Version

Originally published at:

Heimgartner, Heinz; Mlostoń, Grzegorz; Romanski, Jaroslaw (2005). Adamantanethione. In: Paquette, Leo A; Rigby, James H; Crich, David; Wipf, Peter. Electronic Encyclopedia of Reagents in Organic Synthesis (e-EROS). Chichester: John Wiley Sons, RN00504.

DOI: <https://doi.org/10.1002/047084289X.rn00504>

CONTRIBUTION FOR:

ELECTRONIC ENCYCLOPEDIA OF REAGENTS FOR ORGANIC SYNTHESIS

ADAMANTANETHION

Heinz Heimgartner

Organisch-chemisches Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057

Zürich, Switzerland

Grzegorz Mloston, Jaroslaw Romanski

Section of Heteroorganic Compounds, University of Lodz, Narutowicza 68, PL-90-136

Lodz, Poland

CONTACT DETAILS:

Heinz Heimgartner

Organisch-chemisches Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057

Zürich, Switzerland

Phone: +41 1 635 42 82

Fax: +41 1 635 68 36

e-mail: heimgart@oci.unizh.ch

Adamantanthion

Formula (1)

[23695-65-0]

 $C_{10}H_{14}S$

(166.29)

(reagent used as a stable thioketone useful for the preparation of adamantyl substituted products¹ and for studies on the reactivity of thiocarbonyl compounds)

Alternate Name: Tricyclo[3.3.1.1^{3,7}]decan-2-thion

Physical Data: mp 173-174 °C (petroleum ether),² 164-166 °C,³ 160-162 °C (sublimation).⁴

Solubility: soluble in almost all organic solvents; insoluble in water.

Form Supplied in: orange crystals, commercially not available.

Analysis of Reagent Purity: IR,² ¹³C-NMR.⁵

Preparative Methods: the most frequently applied method to prepare the title reagent is the thionation of adamantanone with P₄S₁₀ in pyridine.^{2,6,7} Other thionation reagents such as [(C₆H₁₁)₃Sn]₂S⁸ or (Me₃Si)₂S⁹ as well as H₂S in the presence of *p*-toluene sulfonic acid¹⁰ are also used. Treatment of adamantanone with H₂S/HCl in ethanolic solution at –55 °C affords the hemithioacetal, which decomposes at 100-130 °C/15-100 Torr to yield **1**.²

Purification: column chromatography (silica gel, petroleum ether/dichloromethane); sublimation.^{4,11}

Handling, Storage, and Precautions: camphoraceous odor; stable at low temperature (dry ice) under inert gas, decomposes slowly at room temperature in contact with air; safe in handling.

Conversion to Adamantanone

The reaction of **1** with nitrosonium tetrafluoroborate (NO^+BF_4^-) in dichloromethane at rt leads to adamantanone (**2**) (eq 1).¹² Treatment of **1** with clay-supported cupper nitrate ($\text{Cu}(\text{NO}_3)_2$) under similar conditions gives **2** in 68 % yield.¹³ Formation of **2** (40 %), together with other products, is also reported to occur by ozonolysis in dichloromethane at -70°C .¹⁴

eq (1)

Heating of **1** in ethanol in a sealed tube at 80°C for 14 h yields, after chromatographic workup (SiO_2 , hexane/benzene 10:1), **2** as the major product (67 %) along with di(adamantan-2-yl)-disulfane (20 %).¹⁵ The formation of **2** is also observed when the [2+3] cycloadducts of **1** with nitrile oxides are heated to $185\text{--}250^\circ\text{C}$.¹⁶ Similar treatment of the cycloadduct with diphenyl-nitrile imine leads to adamantanylidene *N*-phenylamine, which can be hydrolyzed in aqueous HCl to give **2**.¹⁶ Hetero-Diels-Alder reaction of **1** with α,β -unsaturated aldehydes and ketones carried out in benzene at 80°C in the presence of 1,4-hydroquinone afford variable amounts of **2**

in addition to the dimer **3** and the corresponding [2+4] cycloadducts as the major products.¹⁵

The latter, being mixed thioacetals, undergo smooth hydrolysis in aqueous ethanolic solution to yield **2**.

Dimerization and Trimerization

Dimerization of **1** leading to 1,3-dithiethane **3** (68 %) occurs smoothly in methane sulfonic acid at rt (eq 2).^{2,7} The dimer **3** is formed as a minor product in attempted hetero-Diels-Alder reactions of α,β -unsaturated ketones with **1** in the presence of hydroquinone at 180 °C.¹⁵ The dimer **3** appears as the major product when a mixture of **1** and 2-(hydroxymethyl)- or 2-[(*N,N*-dimethylamino)methyl]phenols in xylene are heated at 180 °C in the presence of catalytic amounts of 1,4-hydroquinone.¹⁷ Reaction of **1** with PCl_5 in boiling tetrachloromethane leads to the dimer **3**.¹⁸ The same product is formed by direct irradiation ($\lambda = 254 \text{ nm}$ or $> 420 \text{ nm}$) as well as when sensitized with benzophenon.¹⁹

eq (2)

The trimer **4** is obtained from **1** upon treatment with methane sulfonic acid diluted in diethyl ether at rt (eq 2).² Its formation is also reported when a solution of **1** in dichloromethane is stirred in the presence of silica gel.¹⁸ The main product of this reaction is dispiro[1,2,4-trithiolane-3,2':5,2''-diadamantane].

The structures of the dimer **3** and the trimer **4** have been established by X-ray crystallography.¹⁸

Thermal decomposition of **4** at 350-355 °C leads to the monomer **1**.²

Reduction Reactions

Adamantane-2-thiol (**5**) can be conveniently prepared by the reduction of **1** with sodium borohydride in DME solution (eq 3).²⁰ The formation of **5** is also reported to occur in reactions of **1** with organolithium compounds²¹ and with Grignard reagents.²²

eq (3)

Oxidation Reactions

A straightforward procedure for the conversion of **1** in adamantanethione *S*-oxide (adamantanylidene sulfine) (**6**) is using *m*-CPBA in diethyl ether (eq 4).²³ Other procedures to oxidize **1** using dimethyldioxirane in acetone²⁴ or oxone® in a two-phase system¹⁴ are less efficient. Sulfine **6** is not only a superior dipolarophile^{23,25,26} but also reacts with cycloaliphatic thioketones as a 1,3-dipolar species.²⁷

eq (4)

Sulfur Transfer Reactions

In numerous reports, **1** is shown to act as a potent interceptor of sulfur, leading to the reactive adamantanethione *S*-sulfide (adamantanylidene thiosulfine) (**7**).²⁸ This intermediate is believed to exist in equilibrium with the corresponding spirodithiirane **8**. In contrast to the relatively stable **6**, neither **7** nor **8** can be isolated (eq 5).

eq (5)

However, **7** is a reactive 1,3-dipole, which undergoes [2+3] cycloadditions with thiocarbonyl compounds to yield 1,2,4-trithiolane derivatives. For example, heating of **1** and elemental sulfur in xylene leads to the trithiolane **9** (eq 6).²⁹ The same reaction can be performed in toluene/-trichloromethane solution.³⁰ Reaction of **1** with S₈ in the presence of sodium thiophenolate in acetone affords **9** as a minor product, along with 1,2,4,5-tetrathiane (**10**) and small amounts of 1,2,3,5,6-pentathiepane (**11**) (eq 6).³¹ After a reaction time of 7 h, **11** is the major compound of the reaction mixture (78 %). Unexpectedly, **9** is also formed in good yield (69 %) on treatment of a solution of **1** in abs. THF with a catalytic amount of TBAF³² as well as in the presence of silica gel in dichloromethane solution.¹⁸

eq (6)

The formation of **9** is also observed when **1** reacts with aniline in benzene solution at rt. Moreover, in the analogous reaction with *p*-toluene sulfonamide in dioxane, the formation of **9** is accompanied by trimerization and dimerization of **1** to give **4** and **3**.³³

Reactions with Chlorine and Chlorosulfanes

Chlorine adds smoothly to the C=S group of **1** leading to α -chlorosulfanyl chloride **12** (eq 7).³⁴

Analogously, sulfur dichloride converts **1** in the chlorodisulfane (**13**). The reactions of **1** with the stable α -chloro chlorosulfanes (**14**) afford the non-symmetrical di- and trisulfanes (**15**) (eq 7).³⁵

eq (7)

Addition Reactions with Organometal Compounds

Nucleophilic additions of organometal compounds with thiocarbonyl groups are known to occur in a thiophilic or carbophilic fashion.³⁶ The type of the product formed depends on the reagent and on the solvent. For example, prenyl lithium adds to **1** in a carbophilic manner to give the corresponding thiol, which is converted to the stable methyl thioether (**16**) by treatment with methyl iodide.²¹ By contrast, the reaction of **1** with butyllithium in THF gives, after acidic work-up, mainly adamantanethiol (**5**) and small amounts of (adamantan-2-yl)(butyl)sulfane (**17**) as the product of the thiophilic addition.

Formulae (16)-(19)

The addition of methyl magnesium bromide with **1** yields different products dependent on the solvent used: whereas in THF thiophilic addition leads to **18**, the reaction in diethyl ether occurs via the carbophilic addition to give **19**.²² In the reaction of vinyl magnesium bromide, the formation of a thiol resulting from carbophilic addition is reported.²² The same addition mode is observed when propargyl magnesium bromide is used (eq 8). In this case, the initially formed product **20** partially cyclizes to give the spirocyclic dihydrothiophene (**21**).²²

eq (8)

On the other hand, the analogous reaction with allyl magnesium bromide leads to (adamantan-2-yl)(allyl)sulfane exclusively.²² As a plausible interpretation of different results observed with allylic reagents, i.e. carbophilic versus thiophilic addition, a subsequent [2,3] sigmatropic rearrangement is discussed.^{21,37}

Carbophilic additions with **1** are reported for lithiated reagents, which are prepared *in situ* by deprotonation of activated CH compounds with butyllithium in THF. For example, the lithiated (methyl)(*p*-tolyl)sulfone reacts with **1** and, after treatment with methyl iodide followed by elimination of methanethiol, gives an (adamantylidenemethyl)(*p*-tolyl)sulfone.³⁸

Treatment of 2,6-dimethylpyridine (**22**) with butyllithium in THF at $-80\text{ }^{\circ}\text{C}$ and subsequent addition of two equivalents of **1** leads to the dithiol **23** in 65-70 % yield (eq 9).³⁹ In this case, the formation of the new C,C bonds is observed as the exclusive process (carbophilic addition).

eq (9)

Deprotonation of (trimethylsilyl)diazomethane with butyllithium at $-60\text{ }^{\circ}\text{C}$ leads to a carbanionic species (**24**), which adds to **1**. In diethyl ether, two-fold extrusion of sulfur and nitrogen gives the olefinic product (**26**), isolated by distillation (8 %) (eq 10).⁴⁰ On the other hand, the reaction carried out in THF leads to the 1,2,3-thiadiazole derivative **26** (31 %).⁴⁰ The formation of these two products is rationalized by a carbophilic addition of the anion followed by elimination processes leading to **25** or cyclization yielding **26**.

eq (10)

Reactions with Carbenes

Photolysis of a mixture of **1** and diazomethane in an argon matrix at 10 K affords adamantane-thione *S*-methylide (**27**) as the product of the thiophilic attack of methylene (eq 11).⁴¹ The structure of **27** in the matrix is evidenced by spectroscopic methods and by calculations. The ylide **27** undergoes a 1,3-dipolar electrocyclization at ca. 80 K to yield the stable spirothiirane

(28). The formation of other thiocarbonyl ylide derivatives of **1** in photochemical reactions of aromatic diazo compounds, such as diphenyl diazomethane and 9-diazofluorene, are also reported.⁴² Similarly, the oxocarbene generated by photo-rearrangement of benzocyclobutenedione is trapped by **1**.⁴³

eq (11)

Strongly electrophilic dihalocarbenes add easily to **1** leading to the corresponding *gem*-dihalogenated thiiranes. Dichlorocarbene can be generated conveniently either by thermal decomposition of (trichloromethyl)phenyl mercury (Seyferth reagent) or by elimination of hydrogen chloride from trichloromethane in a two-phase system (Makosza reaction).⁴⁴ Unlike other halogenated thiiranes, the corresponding derivative of **1** is a fairly stable compound.

The nucleophilic dimethoxycarbene can be conveniently generated by thermal decomposition of the 1,3,4-oxadiazole derivative (**29**) at 100 °C.⁴⁵ When the decomposition is carried out in toluene in the presence of **1**, dimethoxythiirane (**30**) is produced in 92 % yield (eq 12).⁴⁶ The latter is smoothly desulfurized by treatment with triphenylphosphane to give the ketene acetal (**31**).

eq (12)

The spirocyclic thiirane (**28**) is also available by methylene transfer from triphenylphosphonium methyllide (Wittig reagent) to **1**.⁴⁷ An improved protocol for the isolation of **28**, using carbon disulfide as a solvent and an excess of elemental sulfur, is published.⁴⁸

Formation of Thietanes

The reaction of cycloheptatrienyl carboxylic acid chloride (**32**) and **1** in benzene in the presence of triethylamine gives a mixture of the two products **33** and **34** in 20 and 69 % yield, respectively (eq 13).⁴⁹ The structure of thietane **33** formally corresponds to the [2+2] cycloadduct of **1** and the ketene, which is generated from **32** *in situ*. The formation of **34** is an evidence for a zwitterionic intermediate, in which the ring closure to **34** competes with the cyclization to **33**.

eq (13)

Thietanes (**36**) are formed regioselectively when mixtures of **1** and ketene acetals of type **35** are irradiated in pentane solutions (eq 14).⁵⁰ In the case of the dimethylketene acetal, a mixture of two regioisomeric thietanes is formed.

eq (14)

Irradiation ($\lambda > 420$ nm) of a solution of **1** and α -methylstyrene in benzene affords a mixture of thietane (**38**) and (adamantan-2-yl)(2-phenylallyl)sulfane (**39**) (eq 15).¹⁹ The key intermediate in this reaction is the diradical (**37**), which cyclizes to give **38** or undergoes a 1,5 hydrogen shift to yield **39**.

eq (15)

[2+3] Cycloadditions

Diazo Compounds

Diazomethane reacts with **1** at low temperature to give a mixture of the two regioisomeric [2+3] cycloadducts (**40**) and (**41**) (eq 16).^{51,52} The ratio of the products depends on the polarity of the solvent. Whereas in aprotic, non-polar solvents 2,5-dihydro-1,3,4-thiadiazole (**40**) is the main product, the reaction in methanol results in a 1:9 mixture of **40** and **41**, in favor of the 4,5-dihydro-1,2,3-thiadiazole (**41**).⁵³ Both adducts can be isolated and used for further reactions.^{48,54,55} Thermolysis of **41** in boiling xylene yields thiirane (**28**) as the main product together with homoadamantanethione and small amounts of methylenadamantane. The isomer **40** extrudes nitrogen much easier in a cycloreversion leading to the intermediate adamantanethione *S*-methylide (**27**), which undergoes easily [2+3] cycloadditions with a variety of electron-deficient C,C- and C,X double bonds.⁵⁶⁻⁵⁹

eq (16)

The analogous reaction of **1** with diazoethane affords regioselectively the cycloadduct of type **40**, which can be isolated.^{16,60} The reaction with phenyldiazomethane occurs slower, and at rt spontaneous evolution of nitrogen takes place leading to a phenyl derivative of the thiirane (**28**) via the corresponding thiocarbonyl ylide.¹⁶

Sterically hindered diazocompounds react with **1** to give stable, crystalline cycloadducts, which eliminate nitrogen at elevated temperature to yield thiiranes.^{61,62} By treatment with triphenylphosphane, the latter are converted in sterically congested olefins⁶¹ (two-fold extrusion reaction).⁶³ As an example, the reaction sequence with di(*tert*-butyl)diazomethane is shown in eq 17 (\rightarrow **42** \rightarrow **43** \rightarrow **44**).

eq. (17)

Diphenyldiazomethane and 9-diazofluorene react with **1** in boiling THF to yield the corresponding thiiranes, which are desulfurized in a one-pot procedure to give the aryl-substituted olefins.⁶⁴ A kinetic study of the reaction of **1** with diphenyldiazomethane in DMF at 40 °C is reported.⁶⁵ Ethyl diazoacetate in THF at rt reacts slowly with **1**. The initially formed cycloadduct releases nitrogen spontaneously, and the intermediate thiocarbonyl ylide (**45**) intercepts **1** yielding 1,3-dithiolane (**46**) by a regioselective [2+3] cycloaddition (eq 18).⁶⁶ The reaction is accelerated by

addition of catalytic amounts of lithium perchlorate.⁶⁶ The same product **46** is formed when a solution of dimethyl bromoketene acetal and **1** in pentane is irradiated with light of the wavelength $\lambda > 400$ nm.⁶⁷

eq (18)

The reaction of **1** with α -diazoketones (**47**) in toluene at 90 °C occurs with evolution of nitrogen. The carbonyl-functionalized thiocarbonyl ylide (**48**) undergoes a 1,5-dipolar electrocyclization^{68,69} to give spirocyclic 1,3-oxathiols (**49**) (eq 19).^{70,71}

eq (19)

Similar reactions of **1** with α -diazoamides in THF require catalysis with lithium perchlorate. Depending on the substituent at C(α) of the starting material, the 1,5-dipolar electrocyclization competes with the 1,3-dipolar ring closure to give the thiirane.⁷²

Heating of a three-component mixture consisting of **1**, dimethyl fumarate, and phenylazide in neat leads to the 1,3-oxathiol derivative (**52**), which results from a reaction cascade.⁷³ The first step of this conversion is the [2+3] cycloaddition of phenyl azide to methyl fumarate and subsequent isomerization of dimethyl 4,5-dihydro-1*H*-1,2,3-triazole-4,5-dicarboxylate to the α -diazoester (**50**). Subsequent reaction with **1** and elimination of nitrogen leads to the highly

functionalized thiocarbonyl ylide (**51**), which, after 1,5-dipolar electrocyclization and further transformations, yields (**52**) (eq 20).

eq (20)

Organic Azides

Organic azides react with **1** at 80 °C under evolution of nitrogen and, along with adamantanylidene amines (**56**), 1,2,4-trithiolane (**9**) is always the second product.^{74,75} In these reactions, adamantanethione *S*-imides (**54**), which are generated by nitrogen elimination from the initially formed 1,2,3,4-thiatriazole (**53**), are believed to be key intermediates (eq 21). They undergo ring closure to yield thiaziridines (**55**), which are efficient sulfur donating species by forming imines (**56**). The interception of the sulfur atom by **1** converts it into the reactive thiosulfine (**7**), which undergoes a [2+3] cycloaddition with **1** to give **9**.

eq (21)

In a three-component reaction with **1**, phenyl azide, and fumaronitrile, the formation of **9** is a dominant process, and neither the intermediate *S*-imide (**54**) (R = Ph) nor thiosulfine (**7**) are intercepted.⁷⁶

Nitrile Oxides

Differently substituted nitrile oxides (**57**), which are generated *in situ* by base catalyzed elimination of hydrogen chloride from *N*-hydroxy imidoyl chlorides, in the presence of **1** undergo smoothly [2+3] cycloadditions to yield 1,4,2-oxathiazole derivatives in a regioselective manner.^{16,77} An example with benzonitrile oxide is presented in eq 22.

eq (22)

The cycloadducts of type **58** decompose while heated to 185-250 °C to give the corresponding isothiocyanates and adamantanone (**2**).

Kinetic studies on reactions of substituted adamantanethiones with nitrile oxides are reported.⁷⁸

Nitrile Imides

Similar to the reactions with nitrile oxides, nitrile imides (**59**) undergo regioselective [2+3] cycloadditions with **1** yielding spirocyclic 4,5-dihydro-1,3,4-thiadiazoles (**60**) (eq 23).^{16,77}

Thermal decomposition of these cycloadducts at 200 °C leads to (adamantanylidene)(phenyl) amine and aryl nitrile.

eq (23)

Nitrile Sulfides

Aromatic nitrile sulfides (**62**), which are generated by thermal decarboxylation of 1,3,4-oxathiazol-2-ones (**61**) in boiling xylene, can be trapped by **1** to give the 1,4,2-dithiazole derivatives (**63**) in moderate yields (eq 24).²⁹ The reaction is accompanied by sulfur transfer to **1** and subsequent formation of 1,2,4-trithiolane **9**.

eq (24)

Nitrones

Cyclic and acyclic nitrones are reported to undergo [2+3] cycloadditions with **1**. Typically, the reactions are carried out in toluene or trichloromethane solutions at rt.^{3,79,80} For example, nitrone (**64**) and **1** react to give **65** in 74 % yield (eq 25).⁸⁰ When dissolved in CDCl₃, **65** dissociates to **1** and **64**, and an equilibrium is established proved by ¹H-NMR spectroscopy.⁸⁰

eq (25)

Azomethine Ylides

Thermal ring opening of *N*-substituted aziridines is a convenient method for the generation of azomethine ylides in a stereoselective manner.⁸¹ For example, dimethyl 1,3-diphenylaziridine-

2,2-dicarboxylate (**66**) in boiling toluene, is in equilibrium with azomethine ylide (**67**), which can be intercepted by **1** in a regioselective [2+3] cycloaddition to give **68** in 82 % yield (eq 26).⁸²⁻⁸⁴

eq (26)

Fluoride catalyzed desilylation of properly substituted (trimethylsilyl)methyl amines or (trimethylsilyl)methyl ammonium salts is a superior method for *in situ* generation of azomethine ylides.⁸¹ The polycyclic adduct (**70**) is formed in 87 % yield when **69** is desilylated in the presence of **1** in dichloromethane solution (eq 27).⁸⁵

eq (27)

1,3-Thiazolidines with unsubstituted C(2) and C(4) positions can be obtained from **1** and azomethine ylides without substituents at carbon atoms.^{86,87}

Thiocarbonyl Ylides

Thermal decomposition of 2,5-dihydro-1,3,4-thiadiazoles is a convenient procedure for the generation of thiocarbonyl ylides under mild conditions (−30 to 45 °C).^{58,59} Cycloaliphatic precursors (**71**) heated to 45 °C eliminate nitrogen and the illusive thiocarbonyl ylides (**72**) react with **1** to give regioselectively 2,5-dispiro-1,3-dithiolanes **73** (eq 28).^{55,57,88,89}

eq (28)

The [2+3] cycloaddition of **1** with thiobenzophenone *S*-methylide can be performed at $-30\text{ }^{\circ}\text{C}$ and, in contrast to the above mentioned reactions with **72**, the sterically more hindered regioisomer (**73**) is formed, accompanied by small amounts of 4,4,5,5-tetraphenyl-1,3-dithiolane (eq 29).⁹⁰ It is worth mentioning that the reaction of thiobenzophenone with **27** (generated from **40**) furnishes a mixture of almost equal amounts of the two regioisomeric cycloadducts (**73**) and (**74**).

eq (29)

Thiocarbonyl S-sulfides

Reactions of **1** with thiocarbonyl *S*-sulfides yield 1,2,4-trithiolanes, and the presence of the spiroadamantane residue enhance the stability of these heterocycles significantly. For example, heating of a mixture of **1** and the thermolabile 2,2,5,5-tetraphenyl-1,2,4-trithiolane (**75**) in trichloromethane leads to the mixed trithiolane **76** in 81 % yield (eq 30).^{91,92} An analogous conversion starting with 3,5-di(*tert*-butyl)-3,5-bis(4-methylphenyl)-1,2,4-trithiolane and **1** is also known.⁹³

eq (30)

Three-membered sulfur heterocycles such as thiiranes and *in situ* formed thiaziridines are known as efficient sulfur donors. On the other hand, **1** is recognized as an excellent sulfur interceptor. Therefore, adamantanethione *S*-sulfide (**7**) is frequently postulated as an intermediate involved in the formation of 1,2,4-trithiolanes. The sulfurization of **1** with 2,2-diphenylthiirane yields the trithiolane-bis-spiroadamantane (**9**).⁹²

Thermal three-component reactions with **1**, phenyl azide, and aromatic thioketones furnish mixtures of **9** and mixed 1,2,4-trithiolanes of type **76**.⁷⁴ These reactions are initiated by the [2+3] cycloaddition of phenylazide with the aromatic thioketone, being more reactive than **1**.

The thiaziridine, which is formed after elimination of nitrogen, is considered as the sulfur donor according to the reaction sequence shown in eq (21).

The thermal stability of 1,2,4-trithiolanes containing a spiro-adamantane residue is confirmed by the experiment presented in eq 31.⁹⁴ The initial step of the reaction sequence is the formation of thiobenzophenone from (diphenylmethylidene)triphenylphosphorane and elemental sulfur at 110 °C.

eq (31)

Thiocarbonyl S-oxides

Cycloaliphatic thioketone *S*-oxides (sulfines) (**77**) react with **1** yielding 1,2,4-oxadithiolanes (**78**) as products of a [2+3] cycloaddition (eq 32).⁹⁵ These are the only examples in which sulfines act as 1,3-dipoles. On the one hand, the very high dipolarophilicity of thioketones ('super dipolarophiles') and, on the other hand, steric congestion are of crucial importance for the successful synthesis of **78**, which are obtained in almost quantitative yield.

eq (32)

Thiocarbonyl S-imides

Thiocarbonyl *S*-imides formed as intermediates in reactions of **1** with organic azides do not intercept **1** in the expected [2+3] cycloaddition,^{74,75} in contrast to 2,2,4,4-tetramethyl-3-thioxocyclobutanone.⁹⁶ There is only one report of the reaction of **1** with the stable hexafluorothioacetone *S*-(adamantan-1-yl)imid (**79**).⁹⁷ In this reaction, carried out in deuteriochloro-methane at 100 °C in a sealed tube, the trithiolane **9** is formed as the major product, and 1,4,2-dithiazolidine (**80**), an isomer of the expected [2+3] cycloadduct of **1** and **79**, is obtained in 8 % yield (eq 33). The formation of **80** can be explained by an isomerization of **79** to the corresponding thionitrone *via* a thiaziridine as the intermediate. This interpretation is supported by the formation of **9** in high yield.

eq (33)

Carbonyl Ylides

Heating of 3-phenyloxirane-2,2-dicarbonitrile (**81**) in toluene (110 °C) results in the ring opening leading to the reactive carbonyl ylide (**82**). The latter can be intercepted by **1** to give the 1,3-oxathiolane (**83**). The isomeric product (**84**) results from the nucleophilic ring opening of **81** by the sulfur atom of **1** and subsequent ring closure (eq 34).⁹⁸

eq (34)

Carbonyl O-oxides

The ozonolysis of alkyl vinyl ethers of type **85** in diethyl ether or dichloromethane at –70 °C is an efficient method for *in situ* generation of carbonyl *O*-oxides (**86**), which belong to less explored 1,3-dipolar species. When the reaction is carried out in the presence of **1**, 1,2,4-dioxathiolanes (**87**) are obtained in up to 70 % yield (eq 35).^{14,99} The [2+3] cycloaddition occurs regio-selectively.

eq (35)

Reactions with Oxiranes

Monosubstituted oxiranes undergo a *Lewis*-acid catalyzed reaction with **1** to give 1,3-oxathiolanes in moderate yield.¹⁰⁰ In the presence of zinc chloride in dichloromethane at $-30\text{ }^{\circ}\text{C}$, the reaction with phenyloxirane occurs regioselectively to give the 4-phenyl-1,3-oxathiolane (**89**) (eq 36). Analogous reactions catalyzed with silica gel are performed with methyl- and phenyloxirane leading to mixtures of regioisomeric 1,3-oxathiolanes, e.g. **89** and **90**. Using enantiomerically pure oxiranes, optically active products are obtained.

eq (36)

[2+4] Cycloadditions

Thermal cycloaddition reactions of **1** with conjugated, acyclic and cyclic dienes (hetero *Diels-Alder* reaction) in toluene at variable temperatures in the presence of 1,4-hydroquinone afford thiines of type **92** (eq 37).^{77,101} For example, the reaction with *Danishefsky's* diene (**91**) leads regioselectively to **92**, which, after desilylation, gives **93** in good yield. In the reaction with isoprene, two regioisomeric cycloadducts are formed in comparable amounts.

eq (37)

In analogous reactions, α,β -unsaturated carbonyl compounds and **1** at $80\text{--}140\text{ }^{\circ}\text{C}$ give the expected 1,3-oxathiin along with small amounts of the dimer (**3**) and up to 38 % of **2**.^{15,77}

A special case of heterodienes are *ortho*-quinone methanides of type **96**, which frequently are generated *in situ* by thermal 1,4-dehydration or 1,4-deamination of 2-(hydroxymethyl)phenols and 2-[(*N,N*-dialkylamino)methyl]phenols, respectively. When the precursors **94** or **95** are heated in xylene at 180 °C in the presence of **1** and catalytic amounts of 1,4-hydroquinone, the fused heterocycle **97** is formed (eq 38).^{17,77}

eq (38)

The thermal ring opening of benzothiete leads to the sulfur analogue of **96**, which can be trapped by **1** to give the corresponding 1,3-benzothiin, i.e. the sulfur analogue of **97**.¹⁰²

The formation of 1,3-oxathiin derivatives (**98**) and (**99**) is reported to occur when **1** reacts with acetylene carboxylic acids¹⁰³ or bis(pivaloyl)ketene.¹⁰⁴ The structures of the products are shown in eq 39.

eq (39)

Photochemical Reactions

Irradiation ($\lambda = 470\text{-}500\text{ nm}$) of a solution of **1** in benzene in the presence of equimolar amounts of adamantane-2-thiol (**5**) leads to the formation of the symmetrical di(adamantan-2-yl)disulfane.¹⁰⁵ When **1** is irradiated in hydrocarbons such as cyclohexane, neopentane, butane,

1,1,2,2-tetramethylcyclopropane, etc. with UV light ($\lambda = 254$ nm), mixtures of two isomeric adducts are formed, always accompanied by dimer **3**. A typical example leading to **100** and **101** is shown in eq 40.¹⁰⁶ These conversions are considered to occur *via* singlet π, π^* hydrogen abstraction leading to a radical pair.

eq (40)

Another example with trimethylsilane is described to give the silylated product (**102**) (eq 41).¹⁰⁷

eq (41)

The absorption, emission, and excitation spectra of **1** are described in detail and compared with those of other thioketones.¹⁰⁸

Miscellaneous

Recently, the reaction of **1** with Lawesson's reagent in refluxing deuteriotrichloromethane has been reported to yield spiro-1,3,2-dithiaphosphetane-2-sulfide (**103**) in 87 % yield (eq 42).¹⁰⁹

The structure of this formal [2+2] cycloadduct, which is a sulfur analogue of the postulated intermediates in the thionation of ketones with Lawesson's reagent, has been established by X-ray crystallography.

eq (42)

When heated above the melting point (163-165 °C), the colorless crystals turn to a purple oil indicating the decomposition to **1**.

Heating of a mixture of **1** and 2,3-benzo-7,7-dimethyl-1,4,5,6-tetraphenyl-7-germanonorbornadien (**104**), the precursor of dimethylgermylene, in benzene at 70 °C leads to the 1,3,2,4-dithiadigermolane (**105**) in 38 % yield, together with adamantylideneadamantane (**106**) and 1,2,3,4-tetraphenylnaphthalene (**107**) (eq 43).¹¹⁰ A reaction mechanism via the formation of an intermediate thiagermirane is proposed.

eq (43)

The analogous three-membered intermediate can be trapped in the case of the 7,7-diphenyl-7-germanonorbornadien: after heating of a mixture with **1** in benzene to 80 °C, followed by treatment of the crude product with ethanol at rt, the 2-[(ethoxy)diphenylgermyl]adamantane-2-thiol is obtained in 50 % yield. The latter is believed to be the adduct of ethanol to the thiagermirane.

A thermally more stable thiagermirane (**109**) bearing bulky substituents can be isolated from the photolysis of a mixture of hexamethyltrigermene (**108**) and **1** in cyclohexane using a low-

pressure mercury lamp.¹¹⁰ In addition of 53 % of **109**, 11 % of 1,2,3-thiadigermetane (**110**) and 37 % of the adamantanethione dimer (**3**) are obtained (eq 44). The photolysis and thermolysis of **109** leads to **1** and dimesitylgermylene, which can be intercepted by 2,3-dimethylbuta-1,3-diene. The photolysis of **109** in a hydrocarbon matrix at 77 K leads to the formation of the corresponding thiocarbonyl germide.¹¹⁰

eq (44)

A reduction of **1** to the mercuric thiolate (**112**) is achieved when a solution of **1** in acetonitrile is treated with bis(trifluoroacetato)-1,2-phenylenedimercury (**111**) in the presence of 1-benzyl-1,4-dihydronicotinamide (eq 45).¹¹¹

eq (45)

References

1. Litvinov, V.P., *Chem.Heterocycl. Compounds* **2002**, 38, 9-34.
2. Greidanus, J.W., *Can. J. Chem.* **1970**, 48, 3530-3536.
3. Black, D.S.C.; Watson, K.G., *Aust. J. Chem.* **1973**, 26, 2491-2504.
4. Huisgen, R.; Mloston, G., *Pol. J. Chem.* **1999**, 73, 635-644.
5. Andrieu, C.G.; Debruyne, D.; Paquer, D., *Org. Magn. Reson.* **1978**, 11, 528-532.

6. Falk, K.; Steer, R.P., *Can. J. Chem.* **1988**, *66*, 575-577.
7. Ishii, A.; Akazawa, T.; Ding, M.-X.; Honjo, T.; Maruta, T.; Nakamura, S.; Nagaya, H.; Ogura, M.; Teramoto, K.; Shiro, M.; Hoshino, M.; Nakayama, J., *Bull. Chem. Soc. Jpn.* **1997**, *70*, 509-524.
8. Steliou, K.; Mrani, M., *J. Am. Chem. Soc.* **1982**, *104*, 3104-3106.
9. Degl'Innocenti, A.; Capperucci, A.; Mordini, A.; Reginato, G.; Ricci, A.; Cerreta, F., *Tetrahedron Lett.* **1993**, *34*, 873-876.
10. Kabe, Y.; Takata, T.; Ueno, K.; Ando, W., *J. Am. Chem. Soc.* **1984**, *106*, 8174-8180.
11. Lawrence, A.H.; Liao, C.C.; de Mayo, P.; Ramamurthy, V., *J. Am. Chem. Soc.* **1976**, *98*, 2219-2226.
12. Olah, G.A.; Arvanaghi, M.; Ohannesian, L.; Prakash, G.K.S., *Synthesis* **1984**, 785-786.
13. Chalais, S.; Cornelius, A.; Laszlo, P.; Mathy, A., *Tetrahedron Lett.* **1985**, *26*, 2327-2328.
14. Tabuchi, T.; Nojima, M.; Kusabayashi, S., *J. Chem. Soc., Perkin Trans. I* **1991**, 3043-3046.
15. Katada, T.; Eguchi, S.; Esaki, T.; Sasaki, T., *J. Chem. Soc., Perkin Trans. I* **1984**, 1869-1874.
16. Katada, T.; Eguchi, S.; Sasaki, T., *J. Chem. Soc., Perkin Trans. I* **1984**, 2641-2647.
17. Katada, T.; Eguchi, S.; Esaki, T.; Sasaki, T., *J. Chem. Soc., Perkin Trans. I* **1984**, 2649-2653.
18. Linden, A.; Fu, C.; Majchrzak, A.; Mloston, G.; Heimgartner, H., *Acta Crystallogr., Sect. C* **2002**, *58*, o231-o234.

19. Liao, C.C.; de Mayo, P., *Chem. Commun.* **1971**, 1525-1526.
20. Greidanus, J.W., *Can. J. Chem.* **1970**, *48*, 3593-3597.
21. Rautenstrauch, V., *Helv. Chim. Acta* **1974**, *57*, 496-508.
22. Paquer, D.; Vazeux, M., *J. Organomet. Chem.* **1977**, *140*, 257-264.
23. Zwanenburg, B.; Wagenaar, A.; Thijs, L.; Strating, J., *J. Chem. Soc., Perkin Trans. I* **1973**, 73-75.
24. Tabuchi, T.; Nojima, M.; Kusabayashi, S., *J. Chem. Soc., Chem. Commun.* **1990**, 625-627.
25. Bonini, B.F.; Maccagnani, G.; Mazzanti, G.; Thijs, L.; Ambrosius, H.P.M.M.; Zwanenburg, B., *J. Chem. Soc., Perkin Trans. I* **1977**, 1468-1471.
26. Bonini, B.F.; Maccagnani, G.; Mazzanti, G.; Thijs, L.; Veenstra, G.E.; Zwanenburg, B., *J. Chem. Soc., Perkin Trans. I* **1978**, 1218-1224.
27. Huisgen, R.; Mloston, G.; Polborn, K.; Sustmann, R.; Sicking, W., *Liebigs Ann./Recueil* **1997**, 179-185.
28. Fabian, J.; Senning, A., *Sulfur Rep.*, **1998**, *21*, 1-42.
29. Wai, K.-F.; Sammes, M.P., *J. Chem. Soc., Perkin Trans. I* **1991**, 183-187.
30. Okuma, K.; Kojima, K.; Shibata, S., *Heterocycles* **2000**, *53*, 2753-2758.
31. Huisgen, R.; Rapp, J., *Heterocycles* **1997**, *45*, 507-526.
32. Mloston, G.; Prakash, G.K.S.; Olah, G.A.; Heimgartner, H., *Helv. Chim. Acta* **2002**, *85*, 1644-1658.
33. Campbell, M.M.; Evgenios, D.M., *J. Chem. Soc., Perkin Trans. I* **1973**, 2862-2866.

34. Still, I.W.J.; Kutney, G.W.; McLean, D., *J. Org. Chem.* **1982**, *47*, 555-560.
35. Mloston, G.; Majchrzak, A.; Senning, A.; Sotofte, I., *J. Org. Chem.* **2002**, *67*, 5690-5695.
36. Schaumann, E., In *The Chemistry of Double-bonded Functional Groups*, Vol. 2, Part 2, Supplement A; Patai, S.; Wiley: New York, 1989, pp. 1269-1367.
37. Masson, S.; Saquet, M.; Thuillier, A., *Tetrahedron* **1977**, *33*, 2949-2954.
38. Looutjes, J.A.; Leij, M.; Zwanenburg, B., *Recl. Trav. Chim. Pays-Bas* **1980**, *99*, 39-44.
39. Koning, B.; Hulst, A.; Bouter, A.; Buter, J.; Meetsma, A.; Kellogg, R.M., *Chem. Commun.* **1997**, 1065-1066.
40. Shioiri, T.; Iwamoto, Y.; Aoyama, T., *Heterocycles* **1987**, *26*, 1467-1470.
41. Mloston, G.; Romanski, J.; Schmidt, C.; Reisenauer, H.P.; Maier, G., *Chem. Ber.* **1994**, *127*, 2527-2530.
42. McGimpsy, W.G.; Scaiano, J.C., *Tetrahedron Lett.* **1986**, *27*, 547-550.
43. Boate, D.A.; Johnston, L.J.; Kwong, P.C.; Lee-Ruff, E.; Scaiano, J.C., *J. Am. Chem. Soc.* **1990**, *112*, 8858-8863.
44. Mloston, G.; Romanski, J.; Swiatek, A.; Heimgartner, H., *Helv. Chim. Acta* **1999**, *82*, 946-956.
45. Warkentin, J., *Adv. Carbene Chem.* **1998**, *2*, 245-295.
46. Dawid, M.; Mloston, G.; Warkentin, J., *Chem. Eur. J.* **2002**, *8*, 2184-2187.
47. Krapcho, A.P.; Silvon, M.P.; Flanders, S.D., *Tetrahedron Lett.* **1974**, *15*, 3817-3820.
48. Huisgen, R.; Mloston, G., *Pol. J. Chem.* **1999**, *73*, 635-644.

49. Adam, W.; Peters, E.-M.; Rebollo, H.; Rosenthal, R.J.; Schnering, H.G., *Chem. Ber.* **1984**, *117*, 2393-2408.
50. Ooms, P.; Hartmann, W., *Tetrahedron Lett.* **1987**, *28*, 2701-2704.
51. Krapcho, A.P.; Rao, D.R.; Silvon, M.P.; Abegaz, B., *J. Org. Chem.* **1971**, *36*, 3885-3890.
52. Krapcho, A.P.; Silvon, M.P.; Goldberg, I.; Jahngen Jr., E.G.E., *J. Org. Chem.* **1974**, *39*, 860-861.
53. Sustmann, R.; Sicking, W.; Huisgen, R., *J. Am. Chem. Soc.* **2003**, *125*, 14425-14434.
54. Huisgen, R.; Mloston, G., *Tetrahedron Lett.* **1985**, *26*, 1049-1052.
55. Mloston, G.; Huisgen, R., *Heterocycles* **1985**, *23*, 2201-2206.
56. Mloston, G.; Huisgen, R.; Huber, H.; Stephenson, D.S., *J. Heterocycl. Chem.* **1999**, *36*, 959-968.
57. Mloston, G.; Huisgen, R.; Polborn, K., *Tetrahedron* **1999**, *55*, 11475-11494.
58. Mloston, G.; Heimgartner, H., *Pol. J. Chem.* **2000**, *74*, 1503-1532.
59. Mloston, G.; Heimgartner, H., In *The Chemistry of Heterocyclic Compounds*, Vol. 59: *Synthetic Applications of 1,3-Dipolar Cycloaddition Chemistry Toward Heterocycles and Natural Products*; Padwa, A.; Pearson, W.H., Eds.; Wiley: New York, 2002, pp. 315-360.
60. Huisgen, R.; Li, X.; Mloston, G.; Knorr, R.; Huber, H.; Stephenson, D.S., *Tetrahedron* **1999**, *55*, 12783-12798.
61. Cordt, F.; Frank, R.M.; Lenoir, D., *Tetrahedron Lett.* **1979**, 505-508.
62. Kabe, Y.; Takata, T.; Ueno, K.; Ando, W., *J. Am. Chem. Soc.* **1984**, *106*, 8174-8180.

63. Guziec, F.S.; Sanfilippo, L.J., *Tetrahedron* **1988**, *44*, 6241-6285.
64. Adam, W.; Arias Encarnacion, L.A., *Chem. Ber.* **1982**, *115*, 2592-2605.
65. Huisgen, R.; Langhals, E., *Tetrahedron Lett.* **1989**, *30*, 5369-5372.
66. Kägi, M.; Mloston, G.; Heimgartner, H., *Pol. J. Chem.* **1998**, *72*, 678-687.
67. Ooms, P.H.J.; Hartmann, W., *Angew. Chem., Int. Ed.* **1985**, *24*, 976-977.
68. Taylor, E.C.; Turchi, I.J., *Chem. Rev.* **1979**, *79*, 181-231.
69. Huisgen, R., *Angew. Chem., Int. Ed.* **1980**, *19*, 947-973.
70. Kägi, M.; Linden, A.; Mloston, G.; Heimgartner, H., *Helv. Chim. Acta* **1996**, *79*, 855-874.
71. Kelmendi, B.; Mloston, G.; Heimgartner, H., *Heterocycles* **2000**, *52*, 475-482.
72. Kägi, M.; Linden, A.; Mloston, G.; Heimgartner, H., *Helv. Chim. Acta* **1998**, *81*, 285-302.
73. Mloston, G.; Romanski, J.; Linden, A.; Heimgartner, H., *Pol. J. Chem.* **1996**, *70*, 595-602.
74. Mloston, G.; Romanski, J.; Heimgartner, H., *Pol. J. Chem.* **1996**, *70*, 437-445.
75. Mloston, G.; Romanski, J.; Linden, A.; Heimgartner, H., *Pol. J. Chem.* **1996**, *70*, 880-890.
76. Mloston, G.; Romanski, J.; Linden, A.; Heimgartner, H., *Helv. Chim. Acta* **1996**, *79*, 1305-1314.
77. Sasaki, T.; Eguchi, S.; Katada, T., *Heterocycles* **1984**, *21*, 682.
78. Tsai, T.-L.; Chen, W.-C.; Yu, C.-H.; le Noble, W.J.; Chung, W.-S., *J. Org. Chem.* **1999**, *64*, 1099-1107.
79. Huisgen, R.; Fisera, L.; Giera, H.; Sustmann, R., *J. Am. Chem. Soc.* **1995**, *117*, 9671-9678.
80. Giera, H.; Huisgen, R., *Recl./Liebigs Ann.* **1997**, 1685-1690.

81. Urbaniak, K.; Szymanski, R.; Romanski, J.; Mloston, G.; Domagala, M.; Heimgartner, H., *Helv. Chim. Acta* **2004**, *87*, 496-510.
82. Mloston, G.; Skrzypek, Z., *Bull. Soc. Chem. Belg.* **1990**, *99*, 167-170.
83. Mloston, G.; Urbaniak, K.; Heimgartner, H., *Helv. Chim. Acta* **2002**, *85*, 2056-2064.
84. Domagala, M.; Linden, A.; Olszak, T.A.; Mloston, G.; Heimgartner, H., *Acta Crystallogr., Sect. C.* **2003**, *59*, o250-o253.
85. Butler, R.N.; Farrell, D.M.; McArdle, P.; Cunningham, D., *J. Chem. Soc., Perkin Trans. I* **1998**, 869-873.
86. Gebert, A.; Linden, A.; Mloston, G.; Heimgartner, H., *Pol. J. Chem.* **2003**, *77*, 867-879.
87. Beugelmans, R.; Chastanet, J.; Roussi, G., *Heterocycles* **1987**, *26*, 3197-3202.
88. Huisgen, R.; Mloston, G.; Fulka, C., *Heterocycles* **1985**, *23*, 2207-2212.
89. Huisgen, R.; Mloston, G.; Polborn, K.; Sustmann, R., *Chem. Eur. J.* **2003**, *9*, 2256-2263.
90. Huisgen, R.; Li, X.; Mloston, G.; Fulka, C., *Eur. J. Org. Chem.* **2000**, 1695-1702.
91. Huisgen, R.; Rapp, J., *J. Am. Chem. Soc.* **1987**, *109*, 902-903.
92. Huisgen, R.; Rapp, J., *Tetrahedron* **1997**, *53*, 939-960.
93. Okuma, K.; Shibata, S.; Shioji, K.; Yokomari, Y., *Chem. Commun.* **2000**, 1535-1536.
94. Okuma, K.; Kentaro, K.; Shibata, S., *Heterocycles* **2000**, *53*, 2753-2758.
95. Huisgen, R.; Mloston, G.; Polborn, K.; Sustmann, R.; Sicking, W., *Recl./Liebigs Ann.* **1997**, 179-186.

96. Mloston, G.; Romanski, J.; Linden, A.; Heimgartner, H., *Helv. Chim. Acta* **1993**, 76, 2147-2154.
97. Mloston, G.; Celeda, M.; Roesky, H.W.; Parisini, E.; Ahlemann, J.-T., *Eur. J. Org. Chem.* **1998**, 456-466.
98. Meier, K.-R.; Linden, A.; Mloston, G.; Heimgartner, H., *Helv. Chim. Acta* **1997**, 80, 1190-1204.
99. Tabuchi, T.; Toshihiko, N.; Nojima, M.; Kusabayashi, S., *J. Chem. Soc., Chem. Commun.* **1990**, 625-627.
100. Fu, C.; Linden, A.; Heimgartner, H., *Heterocycles* **2002**, 58, 333-346.
101. Katada, T.; Eguchi, S.; Sasaki, T., *J. Org. Chem.* **1986**, 51, 314-320.
102. Meier, H.; Groschl, D.; Beckert, R.; Weiss, D., *Recl./Liebigs Ann.* **1997**, 1603-1606.
103. Okuma, K.; Maekawa, S.; Shibata, S.; Shioji, K.; Inoue, T.; Kurisaki, T.; Wakita, H.; Yokomari, Y., *Eur. J. Org. Chem.* **2003**, 3727-3729.
104. Kappe, C.O.; Wentrup, C.; Kollenz, G.; *Monatsh. Chem.* **1993**, 124, 1133-1142.
105. Bolton, J.R.; Chen, K.S.; Lawrence, A.H.; de Mayo, P., *J. Am. Chem. Soc.* **1975**, 97, 1832-1837.
106. Law, K.Y.; de Mayo, P., *J. Am. Chem. Soc.* **1979**, 101, 3251-3260.
107. Kira, M.; Akiyama, M.; Ichinose, M.; Sakurai, H., *J. Am. Chem. Soc.* **1989**, 111, 8256-8262.
108. Falk, K.J.; Steer, R.P., *J. Am. Chem. Soc.* **1989**, 111, 6518-6524.

109. Oshida, H.; Ishii, A.; Nakayama, J., *Tetrahedron Lett.* **2004**, *45*, 1331-1334.
110. Tsumuraya, T.; Sato, S.; Ando, W., *Organometallics* **1989**, *8*, 161-167.
111. Wuest, J.D.; Zacharie, B., *J. Am. Chem. Soc.* **1985**, *107*, 6121-6123.